

On the existence periodic orbits in a class of Mechanical Hamiltonian Systems – An elementary mathematical analysis

Luiz C.L. Botelho

Departamento de Matemática Aplicada,
Instituto de Matemática, Universidade Federal Fluminense,
Rua Mario Santos Braga
24220-140, Niterói, Rio de Janeiro, Brazil

Abstract

We present an illustrative application of the two famous Mathematical Theorems in Manifolds in R^N , namely: The Hadamard Theorem on the characterization of convex hypersurfaces in R^N and the well-known Brower fixed point Theorem – in order to show the existence of periodic orbits with an arbitrary given period for a class of Hamiltonian systems. This mathematical result sheds some light on the old mathematical question of existence of periodic orbits in the N -body problem in Classical Mechanics.

1 Elementary may be deep - T. Kato

One of the most fascinating study in Classical – Astronomical Mechanics is that one related to the inquiry on how energetic-positions configurations in a given mechanical dynamical system can lead to periodic systems trajectories for any given period T , a question of basic importance on the problem of the existence of planetary systems around stars in Dynamical Astronomy ([1]).

In this short note we intend to given a mathematical characterization of a class of N -particle Hamiltonian systems possessing intrinsically the above mentioned property of periodic trajectories for any given period T , a result which is obtained by an elementary and illustrative application of the famous Brower fixed point theorem.

As a consequence of this mathematical result we conjecture that the solution of existence of periodic orbits in Classical Mechanics of N -body systems moving through gravitation should be searched with the present status of our Mathematical Knowlegdment.

Let us, thus, start our note by considering a Hamiltonian function $H(x^i, p^i) \equiv H(z^i) \in$

$C^\infty(R^{6N})$ of a system of N -particles set on motion in R^3 . Let us suppose that the Hessian matrix determinant associated to the given Hamiltonian is a positive-definite matrix in the $(6N - 1)$ -dimensional energy-constant phase space $z^i = (x^i, p^i)$ of our given Mechanical system

$$\left\| \frac{\partial^2 H}{\partial z^k \partial z^s} \right\|_{H(z^s)=E} > 0 \quad (1-a)$$

In the simple case of N -particles with unity mass and a positive two-body interaction with a potential $V(|x_i - x_j|^2)$, such that the set in R^n , $\{x^i \mid V(x^i) < E\}$ is a bounded set for every $E > 0$. Our condition eq.(1-a) takes a form involving only the configurations variables $(\{x^p\}_{p=1,\dots,N}; x^p \in R^3)$. Namely

$$||M_{ks}(\{x^p\})|| > 0 \quad (1-b)$$

with

$$\begin{aligned} M_{ks}(\{x^p\}) = \sum_{i < j}^N [V''(|x_i - x_j|^2) 4(x_i - x_j)^2 (\delta_{is} - \delta_{js})(\delta_{ik} - \delta_{jk}) \\ + 2V'(|x_i - x_j|^2)(\delta_{ik} - \delta_{jk})(\delta_{is} - \delta_{js})] \end{aligned} \quad (1-c)$$

The class of mechanical systems given by eq.(1-b)/eq.(1-c) is such that the energy-constant hypersurface $H(z^s) = E$ has always a positive Gaussian curvature and – as a straightforward consequence of the Hadamard Theorem – such energy-constant hypersurface is a convex set of R^{6N} , besides of obviously being a compact set. As a result of the above made remarks, we can see that our mechanical phase space is a convex-compact set of R^{6N} .

The motion equations of the particles of our given Hamiltonian system is given by

$$\frac{dx^i(t)}{dt} = \frac{\partial H}{\partial p^i}(x^i(t), p^i(t)) \quad (2)$$

$$\frac{dp^i(t)}{dt} = -\frac{\partial H}{\partial x^i}(x^i(t), p^i(t))$$

with the initial conditions belonging to our phase space of constant energy $\mathcal{H}_E = \{(x^i, p^i); \mathcal{H}(x^i, p^i) = E\}$, namely:

$$(x^i(t_0), p^i(t_0)) \in \mathcal{H}_E \quad (3)$$

The existence of periodic solutions of the system of ordinary differential equations eq.(2)–eq.(3); for any given period $T \in R^+$ is a consequence of the fact that the Poincaré recurrent application associated to the (global) solutions eq.(2) of the given mechanical system. Note that $H(x^i(t), p^i(t)) = E$

$$P: \mathcal{H}_E \rightarrow \mathcal{H}_E \quad (4)$$

$$(x^i(t_0), p^i(t_0)) \rightarrow (x^i(T), p^i(T))$$

has always a fixed point as a consequence of the application of the Brower fixed point theorem ([3]). This means that there exists always initials conditions in the phase space eq.(3) that produces a periodic trajectory of any period T in our considered class of mechanical N -particle systems.

As a physical consequence, we can see that the existence of planetary systems around stars may be not a rare physical astronomical event, but just a consequence of the mathematical description of the nature laws and the fundamental theorems of Differential Topology and Geometry in R^N .

References

- [1] C.L. Siegel and J.K. Moser – “Lectures on Celestial Mechanics”, Springer-Verlag, Berlin, Heidelberg, New York (1971).
- [2] Shlomo Sternberg – “Lectures on Differential Geometry”, Chelsea Publishing Company, New York (1983).
- [3] Victor Guillemin and Alan Pollack – “Differential Topology”, Prentice-Hall, Inc., Englewood Cliffs, N. Jersey (1974).